©2001-2005 Mineral Data Publishing, version 1

**Crystal Data:** Orthorhombic. *Point Group:* n.d. As small polycrystalline grains, to 0.2 mm; randomly oriented fibers and wiry aggregates.

**Physical Properties:** Hardness = n.d. VHN = 58 (25 g load). D(meas.) = n.d. D(calc.) = 4.50

**Optical Properties:** Opaque. *Color:* Red; in polished section, pale gray. *Luster:* Metallic.  $R_1-R_2$ : (400) 39.0–42.0, (420) 36.8–39.4, (440) 34.6–36.8, (460) 33.0–34.8, (480) 32.1–33.6, (500) 31.4–32.7, (520) 30.8–32.2, (540) 30.4–31.8, (560) 30.0–31.5, (580) 29.6–31.0, (600) 29.1–30.6, (620) 28.7–30.3, (640) 28.4–30.0, (660) 28.0–29.7, (680) 27.6–29.3, (700) 27.1–29.0

**Cell Data:** Space Group: n.d. a = 3.576(2) b = 6.759(2) c = 10.074(5) Z = 2

X-ray Powder Pattern: Duranus, France.

2.919 (10), 5.620 (9), 5.037 (9), 1.969 (9), 1.788 (9), 2.682 (8), 3.016 (7)

Chemistry:		(1)	(2)	(3)
	$\mathbf{As}$	90.0	90.8	90.33
	$\mathbf{S}$	10.3	10.3	9.67
	Total	100.3	101.1	100.00

(1) Duranus, France; by electron microprobe, corresponding to  $As_{3.95}S_{1.05}$ . (2) Do.; corresponding to  $As_{3.96}S_{1.04}$ . (3)  $As_4S$ .

Occurrence: In calcite veinlets in marls and siliceous limestones (Duranus, France).

Association: Arsenic, realgar, orpiment, stibnite, sphalerite, rhodochrosite, quartz, calcite.

**Distribution:** From Duranus, Alpes-Maritimes, France [TL]. In the Mt. Washington copper mine, Vancouver Island, British Columbia, Canada. At the Capillitas mine, San Luis, Catamarca Province, Argentina.

Name: For the locality at Duranus, France.

Type Material: National School of Mines, Paris, France.

**References:** (1) Johan, Z., C. Laforêt, P. Picot, and J. Ferard (1973) La duranusite,  $As_4S$ , un nouveau minéral. Bull. Soc. fr. Minéral., 96, 131–134 (in French). (2) (1975) Amer. Mineral., 60, 945 (abs. ref. 1).