$\odot$  2001-2005 Mineral Data Publishing, version 1

**Crystal Data:** Monoclinic. *Point Group: m.* Rarely as crystals, flattened on  $\{010\}$ , elongated along [001], showing  $\{110\}$ ,  $\{010\}$ ,  $\{011\}$ ,  $\{\overline{1}12\}$ , several other forms. Commonly acicular, fibrous, to 1.5 cm; may be botryoidal to stalactitic.

**Physical Properties:** Cleavage: On  $\{010\}$ , perfect. Fracture: Uneven. Tenacity: Flexible. Hardness = 2–2.5 D(meas.) = 2.53–2.725 D(calc.) = 2.703

**Optical Properties:** Transparent to translucent. *Color:* Colorless, white, pale gray; colorless in transmitted light. *Luster:* Vitreous, pearly on cleavages. *Optical Class:* Biaxial (–). *Orientation:* Z = b;  $X \wedge c = -29^{\circ}$ . *Dispersion:* r > v.  $\alpha = 1.583$  $\beta = 1.589$   $\gamma = 1.594$  2V(meas.) =  $79^{\circ}24'$ 

**Cell Data:** Space Group: Ia. a = 5.9745(5) b = 15.4340(15) c = 6.2797(6) $\beta = 114^{\circ}50(1)'$  Z = 4

**X-ray Powder Pattern:** Sainte-Marie-aux-Mines, France. 4.30 (10), 7.7 (9), 2.70 (9), 2.66 (9), 3.08 (8), 3.02 (8), 5.1 (7)

Chemistry:		(1)	(2)
	$P_2O_5$	0.30	
	$As_2O_5$	50.54	53.19
	$\mathrm{SiO}_2$	0.70	
	$\mathrm{Fe}_2\mathrm{O}_3$	0.35	
	MgO	0.50	
	CaO	23.90	25.96
	$H_2O$	23.80	20.85
	Total	100.09	100.00
(1) $C \rightarrow M \rightarrow M \rightarrow D \rightarrow D$			

(1) Sainte-Marie-aux-Mines, France. (2)  $Ca(HAsO_4) \cdot 2H_2O$ .

**Occurrence:** An uncommon secondary mineral formed by oxidation of earlier arsenic-bearing minerals.

Association: Picropharmacolite, hörnesite, haidingerite.

**Distribution:** In Germany, from Wittichen, Black Forest, in the Anton and Sofia mines; at St. Andreasberg, Harz Mountains; in the Rosenkrantz mine, Schneeberg, Saxony. At Nowa Ruda, Poland. Long crystals from Jáchymov (Joachimsthal), Czech Republic. At Săcărîmb, Romania. From Sainte-Marie-aux-Mines, Haut-Rhin, France. Found at Bou Azzer, Morocco. In the USA, at the Getchell mine, Potosi district, Humboldt Co., and the White Caps mine, Manhattan district, Nye Co., and the Gold Bar mine, Antelope district, Eureka Co., Nevada; at Sterling Hill, Ogdensburg, Sussex Co., New Jersey; in the OK mine, San Gabriel Canyon, Los Angeles Co., California. Some additional older localities may require modern confirmation.

**Name:** From the Greek for *poison*, as an arsenic-containing mineral.

**References:** (1) Palache, C., H. Berman, and C. Frondel (1951) Dana's system of mineralogy, (7th edition), v. II, 706–708. (2) Pierrot, R. (1964) Contribution à la minéralogie des arséniates calciques et calcomagnésiens naturels. Bull. Soc. fr. Minéral., 87, 169–211 (in French). (3) Ferraris, G. (1969) The crystal structure of pharmacolite, CaH(AsO<sub>4</sub>)•2H<sub>2</sub>O. Acta Cryst., 25, 1544–1550. (4) Ferraris, G., D.W. Jones, and J. Yerkess (1971) Determination of hydrogen atom positions in the crystal structure of pharmacolite, CaHAsO<sub>4</sub>•2H<sub>2</sub>O, by neutron diffraction. Acta Cryst., 27, 349–354.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.