(c)2001-2005 Mineral Data Publishing, version 1

Crystal Data: Orthorhombic. Point Group: 2/m 2/m 2/m. Lathlike crystals, to 5 mm, flattened on {100}, elongated and striated along [010], showing {001}, {100}, {021}, {201}, {401}, with rough {010}, {021} as terminations; compact massive.

Physical Properties: Cleavage: On $\{100\}$, fair; on $\{001\}$, $\{021\}$, poor. Fracture: Even to conchoidal. Tenacity: Brittle. Hardness = 4.5 D(meas.) = 2.935(5) D(calc.) = 2.939

Optical Properties: Transparent. Color: Pale brown; colorless in transmitted light. Optical Class: Biaxial (-). Pleochroism: Moderate; X = very pale brown to colorless; Y = pale amber; Z = yellowish brown. Orientation: X = c; Y = b; Z = a. Dispersion: r < v, strong. Absorption: X > Y > Z. $\alpha = 1.631$ –1.648 $\beta = 1.640$ –1.660 $\gamma = 1.641$ –1.663 $2V(\text{meas.}) = 15^{\circ}$ –28°

Cell Data: Space Group: Pbam. a = 8.27-8.289 b = 13.25-13.357 c = 9.01-9.057 Z = 4

X-ray Powder Pattern: Franklin, New Jersey, USA. 3.974 (100), 2.600 (72), 2.184 (49), 2.128 (37), 2.264 (32), 1.708 (32), 1.641 (29)

Chemistry:	(1)	(2)	(3)		(1)	(2)	(3)
B_2O_3	32.40	30.2	34.15	ZnO	3.13		
SiO_2			1.37	MgO	1.66		7.59
Fe_2O_3			0.13	CaO	25.40		23.96
FeO			1.20	H_2O	8.51	11.9	13.40
MnO	28.30		17.87	insol.	0.84		
				Total	100.24		99.67

(1) Franklin, New Jersey, USA; corresponding to $Ca_2(Mn_{1.67}Mg_{0.16}Zn_{0.06})_{\Sigma=1.89}B_4O_7(OH)_6$. (2) Do.; partial analysis; by electron microprobe, B_2O_3 on a separate sample, H_2O by DTA–TGA;

(2) Do.; partial analysis; by electron microprobe, B_2O_3 on a separate sample, H_2O by DTA–TGA; with cations stated to correspond to $Ca_{1.91}(Mn_{1.89}Mg_{0.16}Zn_{0.04})_{\Sigma=2.09}B_4O_7(OH)_6$. (3) Solongo deposit, Russia; corresponding to $Ca_{1.76}(Mn_{1.04}Mg_{0.78}Fe_{0.06}^{2+})_{\Sigma=1.88}B_{4.04}O_{6.67}(OH)_{6.06}$.

Polymorphism & Series: Forms a series with fedorovskite.

Occurrence: Very rare in hydrothermal veins in a metamorphosed stratiform zinc orebody (Franklin, New Jersey, USA).

Association: Clinochrysotile, calcite, franklinite, willemite, zincite (Franklin, New Jersey, USA).

Distribution: From Franklin, Sussex Co., New Jersey, USA. At the Solongo boron deposit, Buryatia, Russia.

Name: To honor George Rowe (1868–1947), Mine Captain at Franklin, New Jersey, USA, and long-time collector of Franklin minerals.

Type Material: National School of Mines, Paris, France; Harvard University, Cambridge, Massachusetts, 96252; National Museum of Natural History, Washington, D.C., USA, 105493, C6291.

References: (1) Palache, C., H. Berman, and C. Frondel (1951) Dana's system of mineralogy, (7th edition), v. II, 377–378. (2) Aristarain, L.F., R.C. Erd, and G.D. Eberlein (1974) Roweite from Franklin, New Jersey: a restudy. Amer. Mineral., 59, 66–70. (3) Moore, P.B. and T. Araki (1974) Roweite, $Ca_2Mn_2^{2+}(OH)_4[B_4O_7(OH)_2]$: its atomic arrangement. Amer. Mineral., 59, 60–65. (4) Malinko, S.V., T.I. Stolyarova, and D.P. Shashkin (1972) The first finding of magnesium roweite, its paragenesis and products of replacement. Zap. Vses. Mineral. Obshch., 101, 465–473 (in Russian). (5) Yamnova, N.A., M.A. Simonov, and N.V. Belov (1975) Crystallochemical structure of roweite, $Ca_2(Mn, Mg)_2[B_4O_7(OH)_2](OH)_4$. Doklady Acad. Nauk SSSR, 221, 1326–1329 (in Russian). (6) Dunn, P.J. (1995) Franklin and Sterling Hill, New Jersey. No publisher, n.p., 641–642.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.