Crystal Data: Hexagonal. Point Group: $6/m \ 2/m \ 2/m$, $\overline{6}m2$, or 6mm. As flattened, radial clusters of platy crystals, to about 0.1 mm.

Physical Properties: Cleavage: Perfect on $\{0001\}$. Hardness = ~ 3 D(meas.) = 2.85(5)D(calc.) = 2.90

Optical Properties: Semitransparent. Color: Dark red. Streak: Pale orange.

Luster: Vitreous on cleavage surfaces; slightly resinous on fracture surfaces.

Optical Class: Uniaxial (-). Pleochroism: O = reddish orange; E = pale pinkish orange.

Absorption: Moderate; O > E. $\omega = 1.728(4)$ $\epsilon = \text{n.d.}$

Cell Data: Space Group: $P6_3/mmc$, $P\overline{6}2c$, or $P6_3/mc$. a = 6.506(7) c = 23.49(3) Z = [1]

X-ray Powder Pattern: Sterling Hill, New Jersey, USA.

11.5 (100), 5.61 (90), 2.844 (60), 2.748 (50), 2.545 (50), 4.56 (40), 3.25 (40)

Chemistry:

	(1)	(2)
$\mathrm{As_2O_5}$	27.4	30.52
$\overline{\mathrm{SiO}_2}$	1.7	
FeO	0.3	
MnO	27.0	28.26
CuO	3.3	
ZnO	0.0	
MgO	0.9	
CaO	12.4	14.90
$\mathrm{H_2O}$	[27.0]	26.32
Total	100.0	100.00

(1) Sterling Hill, New Jersey, USA; H_2O by difference. (2) $Ca_4Mn_6(AsO_4)_4(OH)_8 \cdot 18H_2O$.

Occurrence: Extremely rare in massive granular franklinite-willemite ore from a metamorphosed stratiform zinc orebody.

Association: Manganoan cuprian adamite, franklinite, willemite, calcite.

Distribution: From Sterling Hill, Ogdensburg, Sussex Co., New Jersey, USA.

Name: For the dell of the Wallkill River, in which both the Sterling Hill and the Franklin deposits were discovered.

Type Material: Harvard University, Cambridge, Massachusetts, 113445; National Museum of Natural History, Washington, D.C., USA, 149767.

References: (1) Dunn, P.J. and D.R. Peacor (1983) Kittatinnyite and wallkilldellite, silicate/arsenate analogues containing calcium and manganese, from Franklin and Sterling Hill, New Jersey. Amer. Mineral., 68, 1029–1032.