


Yukonite

(c)2001-2005 Mineral Data Publishing, version 1

Crystal Data: Amorphous, gellike to very poorly crystalline. *Point Group:* n.d.
As irregular concretionary masses, typically strongly cracked.

Physical Properties: *Fracture:* Smooth to conchoidal. *Tenacity:* Extremely brittle.
Hardness = 2–3 D(meas.) = 2.65; 2.86 after gas evolution. D(calc.) = n.d. May decrepitate when fresh, on exposure to air, H₂O, or warmth, with evolution of primarily CO₂.

Optical Properties: Translucent. *Color:* Dark brown, brownish black, violet to deep blood-red; in thin splinters, deep brown, yellowish brown, reddish purple. *Streak:* Brownish yellow. *Luster:* Vitreous to resinous.
Optical Class: Isotropic. *n* = n.d.

Cell Data: *Space Group:* n.d. *Z* = n.d.

X-ray Powder Pattern: Tagish Lake, Canada.

14.1 (100), 2.79 (60), 3.25 (57), 5.58 (37), 2.61 (20), 1.63 (20), 2.24 (11)

Chemistry:	(1)	(2)	(3)		(1)	(2)	(3)
SO ₃		0.12		ZnO		0.56	
P ₂ O ₅		0.22		MgO		0.41	
As ₂ O ₅	36.3	39.68	38.95	CaO	10.6	12.86	13.30
SiO ₂		0.21		H ₂ O ⁺		[5.22]	
Fe ₂ O ₃	36.6	30.25	32.48	H ₂ O ⁻		[10.03]	
MnO		0.44		H ₂ O	17.9		15.27
				Total	101.4	[100.00]	100.00

(1) Tagish Lake, Canada; by electron microprobe, total Fe as Fe₂O₃, H₂O by TGA.

(2) Rędziny, Poland; by electron microprobe, total Fe as Fe₂O₃, H₂O calculated for charge balance; corresponds to (Ca_{6.48}Mg_{0.29}Zn_{0.19}Mn_{0.17})_{Σ=7.13}Fe_{10.70}[(AsO₄)_{9.75}(SiO₄)_{0.10}(PO₄)_{0.09}(SO₄)_{0.04}]_{Σ=9.98}(OH)_{16.37} • 15.72H₂O. (3) Ca₇Fe₁₂(AsO₄)₁₀(OH)₂₀ • 15H₂O.

Occurrence: A secondary mineral typically altered from arsenopyrite.

Association: Symplectite, argentian galena, pyrrargyrite, argentite, chalcopyrite, arsenopyrite, quartz (Tagish Lake, Canada); parasymplectite, köttigite, ogdensburgite, pharmacosiderite, legrandite, willemite, franklinite, sphalerite (Sterling Hill, New Jersey, USA); arsenopyrite, arseniosiderite, arsenolite, barian pharmacosiderite (Trout Creek, Colorado, USA); arsenopyrite, pharmacosiderite (Rędziny, Poland).

Distribution: Found on the west side of Windy Arm, Tagish Lake, Yukon Territory, Canada. In the USA, from Sterling Hill, Ogdensburg, Sussex Co., New Jersey, and the Crystal No. 8 mine, Trout Creek pegmatites, Chaffee Co., Colorado. In Germany, from Saalfeld, Thuringia; at Graulau mountain, near Lammersdorf, Eifel district; from Hasserode, near Wernigerode, Harz Mountains; and at Heubachtal, Black Forest. From Rędziny, Poland.

Name: For Yukon Territory, Canada, within which the mineral was first found.

Type Material: The Natural History Museum, London, England, 1916,454; Geological Survey of Canada, Ottawa, 18594; Royal Ontario Museum, Toronto, Canada, M11468; National Museum of Natural History, Washington, D.C., USA, R5783.

References: (1) Palache, C., H. Berman, and C. Frondel (1951) Dana's system of mineralogy, (7th edition), v. II, 953–955 [arseniosiderite, part]. (2) Tyrrell, J.B. and R.P.D. Graham (1913) Yukonite, a new hydrous arsenate of iron and calcium, from the Tagish Lake, Yukon Territory, Canada; with a note on the associated symplectite. Trans. Roy. Soc. Canada, 7(IV), 3, 13–18. (3) Dunn, P.J. (1982) New data for pitticite and a second occurrence of yukonite at Sterling Hill, New Jersey. Mineral. Mag., 46, 261–264. (4) Pieczka, A., B. Gołębiowska, and W. Franus (1998) Yukonite, a rare Ca-Fe arsenate, from Rędziny (Sudetes, Poland). Eur. J. Mineral., 10, 1367–1370. (5) Ross, D.R. and J.E. Post (1997) New data on yukonite. Powder Diffraction, 12, 113–116.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.