(c)2001-2005 Mineral Data Publishing, version 1

Crystal Data: Hexagonal. *Point Group:* 6/m or 6. Massive, in veinlets, to 0.2 mm wide, and as granular inclusions.

Physical Properties: Fracture: Uneven. Tenacity: Brittle. Hardness = n.d. VHN = 1030 (40 g load). D(meas.) = n.d. D(calc.) = 7.03 Bright yellow-green cathodoluminescence.

Optical Properties: Transparent. Color: Colorless. Luster: Adamantine. Optical Class: Uniaxial. Anisotropism: Noted. Bireflectance: Noted. R₁-R₂: (486) 18.0-19.0, (553) 18.6-19.5, (589) 18.9-19.6, (656) 18.5-18.7

Cell Data: Space Group: $P6_3/m$ or $P6_3$ (?). a = 6.231 c = 36.77 Z = 6.231

X-ray Powder Pattern: Near the Irtysh River, Kazakhstan. 3.029 (10), 3.082 (9), 2.780 (9), 6.13 (7), 5.18 (6), 2.481 (6), 1.929 (4)

Chemistry:		(1)
	$\mathrm{Ta_2O_5}$	80.00
	Nb_2O_5	12.07
	$\mathrm{Sn}ar{\mathrm{O}}_2$	0.58
	$\overline{\mathrm{MnO}}$	trace
	PbO	0.41
	CaO	0.20
	$\mathrm{Na_2O}$	6.46
	Total	99.72

(1) Near the Irtysh River, Kazakhstan; by electron microprobe, average of three analyses; corresponds to $(Na_{1.83}Ca_{0.03}Pb_{0.02})_{\Sigma=1.88}(Ta_{3.19}Nb_{0.80}Sn_{0.03})_{\Sigma=4.02}O_{11}$.

Occurrence: As veinlets in altered thoreaulite, and inclusions in ixiolite and lithiotantite, in granite pegmatites.

Association: Thoreaulite, ixiolite, lithiotantite.

Distribution: From the Ungursai tantalum deposit, Kalba Mountains, near the Irtysh River, eastern Kazakhstan.

Name: For the Irtysh River, Kazakhstan, near which it occurs.

Type Material: A.E. Fersman Mineralogical Museum, Academy of Sciences, Moscow, Russia, 43776.

References: (1) Voloshin, A.V., Y.A. Pakhomovskii, L.V. Bulgak, and G.A. Perlina (1985) Irtyshite, a new mineral from granite pegmatites. Mineral. Zhurnal, 7(3), 83–87 (in Russian). (2) (1986) Amer. Mineral., 71, 1545 (abs. ref. 1). (3) (1986) Mineral. Abs., 37, 530 (abs. ref. 1).