©2001-2005 Mineral Data Publishing, version 1

Crystal Data: Monoclinic. *Point Group:* 2/m. Crystals commonly equant, $\{100\}$, $\{001\}$, $\{111\}$, $\{11\overline{1}\}$, and a dozen other forms; also tabular on $\{100\}$, to 3 mm; pseudohexagonal.

Physical Properties: Cleavage: Perfect on $\{100\}$. Fracture: Conchoidal. Tenacity: Brittle. Hardness = 1.5-2 VHN = n.d. D(meas.) = 4.88 D(calc.) = 4.926

Optical Properties: Transparent; becoming opaque on exposure to light, then translucent in thin splinters. *Color:* Scarlet, vermilion, pink; becoming red-orange on exposure to light. *Streak:* Vermilion. *Luster:* Adamantine.

Optical Class: Biaxial (-). Pleochroism: Slight in transmitted light. Orientation: Y = b; $Z \wedge c = 6.5^{\circ}$. Dispersion: Strong. n = 3.27(9) 2V(meas.) = $\sim 65^{\circ}$ R₁-R₂: (400) 40.7-44.0, (420) 40.0-43.2, (440) 39.3-42.4, (460) 38.7-41.5, (480) 37.9-40.4, (500) 37.0-39.2, (520) 36.0-38.0, (540) 35.0-36.6, (560) 33.8-35.4, (580) 32.7-34.7, (600) 32.0-34.0, (620)

 $37.0-39.2, (520) \ 36.0-38.0, (540) \ 35.0-36.0, (560) \ 33.8-35.4, (580) \ 32.7-34.7, (600) \ 32.0-34.0, (620) \ 31.4-33.4, (640) \ 31.0-32.9, (660) \ 30.6-32.4, (680) \ 30.4-31.8, (700) \ 30.1-31.4$

Cell Data: Space Group: A2/a. a = 17.23 b = 7.78 c = 15.19 $\beta = 101^{\circ}12'$ Z = 24

X-ray Powder Pattern: Synthetic.

2.82(100), 3.21(80), 2.72(60), 1.953(50), 1.701(40), 1.661(40), 1.608(40)

Chemistry:		(1)	(2)
	Ag	43.9	43.69
	As	28.9	30.34
	Sb	0.4	
	\mathbf{S}	26.0	25.97
	Total	99.2	100.00

(1) Binntal, Switzerland. (2) $AgAsS_2$.

Polymorphism & Series: Dimorphous with trechmannite.

Occurrence: Of hydrothermal origin with other Ag–As sulfides.

Association: Sphalerite, pyrite, realgar, orpiment, hutchinsonite, sartorite, rathite (Binntal, Switzerland); chabournéite, pierrotite, parapierrotite, stibnite, pyrite, sphalerite, twinnite, zinkenite, madocite, andorite, laffittite, routhierite, aktashite, wakabayashilite, realgar, orpiment (Jas Roux, France); proustite, gersdorffite, argentian miargyrite (Silvermines, Ireland).

Distribution: From the Lengenbach quarry, Binntal, Valais, Switzerland [TL]. At the Jas Roux deposit, 10 km east of Chapelle-en-Valgaudemar, Hautes-Alpes, France. From Wiesloch, Black Forest, Germany. At Silvermines, Co. Tipperary, Ireland.

Name: In honor of George Frederick Herbert Smith (1872–1953), crystallographer of the British Museum (Natural History), London, England.

Type Material: n.d.

References: (1) Palache, C., H. Berman, and C. Frondel (1944) Dana's system of mineralogy, (7th edition), v. I, 430–432. (2) Hellner, E. and H. Burzlaff (1964) Die Struktur des Smithits AgAsS₂. Naturwiss., 51, 35–36 (in German). (3) Berry, L.G. and R.M. Thompson (1962) X-ray powder data for the ore minerals. Geol. Soc. Amer. Mem. 85, 142–143.